

Managing Pulmonary Situations

Better than ever

Program Description:

This innovative, reality based program is designed to involve the learner in real life situations regarding the identification and proper management of normal and abnormal pulmonary situations. The learner has three opportunities for learning material, i.e. classes (video with illustrations), library (animated books) and near virtual reality interactive rooms. The content presented is based on the best available science (evidence based practice), using national or international standards when possible. The program is targeted to the practicing clinicians. It is not designed to create experts, but clinicians who are competent at addressing the major aspects of pulmonary monitoring. References are provided for additional learning opportunity. Capnography has the unique ability to aid clinicians in assessing both ventilation and blood flow. In this program, a review of the physiology that allows exhaled CO₂ to monitor ventilation and perfusion is presented. The technology required for measuring exhaled CO₂ is presented in order to allow the attendee to identify which devices are best for their clinical practice. Due to the ability to assess both ventilation and perfusion, capnography has been called the "15 second vital sign". The use of capnography is likely to quickly grow throughout the hospital and even into pre hospital settings.

Using research that indicates how learning best takes place, our program is structured to engage the learner, using real life learning skills. Our learners fill whatever learning needs they have in our short, targeted classroom videos or learn on their own in our interactive library. Once the learner feels ready to proceed, they enter our virtual reality hospital. The clinical areas are divided into learning/practicing rooms and testing/evaluation rooms.

The program is web based so learning can occur anytime, anywhere. The learner can complete the class in a given time period, e.g. 2-4 weeks. The specific clinical area the learner is employed can be completed in about 2 hours, including the testing/evaluation phase. Except for the testing section, the learner can enter and leave the program whenever they want. This allows improved use so the learner can learn at their convenience, either at work or at home.

Program Objective:

The objective of this program is to:

Develop the knowledge base of practicing clinicians to recognize and successfully manage patients with normal and abnormal pulmonary situations.

Participant Learning Objectives:

At the completion of this course, the participant will be able to:

- 1) Recall the relationship between PaCO₂ and PetCO₂ levels
- 2) Discuss how the reason why capnography is a rapid assessment of inadequate ventilation
- 3) Discuss how capnography can reflect a reduction in cardiac output
- 4) Describe the 4 key applications of capnography in patient monitoring
- 5) Discuss how pulse oximetry acts as an early warning sign of pulmonary (intrapulmonary shunting) changes
- 6) Discuss the limitations of oximetry in evaluating ventilation
- 7) Describe the key errors associated with pulse oximetry
- 8) Describe how to interpret the 8 most common blood gas disturbances
- 9) Discuss the management of respiratory and metabolic acidosis
- 10) Discuss the management of respiratory and metabolic alkalosis

Learning Rooms -The learner enters rooms in each clinical area. Each room has a different problem to solve. As the learner enters a room, they must act as they would in real life. The room is interactive, including the patient. If the learner wants information, they must seek the information. As in real life, no information is given without their effort. The learner must make an assessment and if treatment is indicated, administer that treatment. Treatments can be accessed in real life format, e.g. medication carts. Depending on the treatment selected, the patient may get better. Incorrect answers are identified for the learner and suggestions are made to help learning. The learner must solve

the problem before being allowed to advance. However, the learner can leave the room if they want and complete the same room at a later time.

There are 33 learning rooms with 11 for capnography, 10 for pulse oximetry and 12 for blood gas analysis. Each case is designed to teach specific aspects of identifying and treating pulmonary function. The key teaching points in each room are identification of normal and abnormal situations and applying the correct treatment to optimize the patient.

Testing Rooms – in the evaluation section, the hospital room is identical to the learning rooms. However, in these rooms, once the learner has selected a response, no guidelines are given whether the answer is correct or not. The learner chooses to exit the room at any time. Success will be identified by if the learner has completed all the required steps to ensure the patient was adequately treated.

The testing cases involve four of the most important aspects of pulmonary disturbances and treatment, including potential life threatening situations.

Bibliography / Reference List

Capnography

1. Ackerman MH, Mick DJ. Technologic approaches to determining proper placement of enteral feeding tubes. *AACN Adv Crit Care*. 2006 Jul-Sep;17(3):246-9.
2. Ahrens T, Sona C. Capnography application in acute and critical care. *AACN Clinical Issues* 2003;14:123-132.
3. Ahrens TS, Wiejera H, Ray SE. Capnography. *Critical Care Nursing Clinics of North America* 1999;11:49-62.
4. Baraka AS, Aouad MT, Jalbout MI, Kaddoum RN, Khatib MF, Haroun-Bizri ST. End-tidal CO₂ for prediction of cardiac output following weaning from cardiopulmonary bypass. *J Extra Corpor Technol*. 2004 Sep;36(3):255-7.
5. Burns SM, Carpenter R, Blevins C, Bragg S, Marshall M, Browne L, Perkins M, Bagby R, Blackstone K, Truwit JD. Detection of inadvertent airway intubation during gastric tube insertion: Capnography versus a colorimetric carbon dioxide detector. *Am J Crit Care*. 2006 Mar;15(2):188-95.
6. Burton JH, Harrah JD, Germann CA, Dillon DC. Does end-tidal carbon dioxide monitoring detect respiratory events prior to current sedation monitoring practices? *Acad Emerg Med*. 2006 May;13(5):500-4.
7. Corbo J, Bijur P, Lahn M, Gallagher EJ. Concordance between capnography and arterial blood gas measurements of carbon dioxide in acute asthma. *Ann Emerg Med*. 2005 Oct;46(4):323-7.
8. Fu ES, Downs JB, Schweiger JW, Miguel RV, Smith RA. Supplemental oxygen impairs detection of hypoventilation by pulse oximetry. *Chest*. 2004 Nov;126(5):1552-8.
9. Gazmuri RJ, Kube E. Capnography during cardiac resuscitation: a clue on mechanisms and a guide to interventions. *Crit Care* 2003;7(6):411-412.
10. Grmec S. Comparison of three different methods to confirm tracheal tube placement in emergency intubation. *Intensive Care Med*. 2002 Jun;28(6):701-4.
11. Kline JA, Arunachlam M. Preliminary study of the capnogram waveform area to screen for pulmonary embolism. *Ann Emerg Med*. 1998 Sep;32(3 Pt 1):289-96.
12. Krauss B, Heightman AJ. 15-second triage tool. The use of capnography for the rapid assessment & triage of critically injured patients & victims of chemical terrorism. *JEMS*. 2006 Jun;31(6):60-2, 64-6, 68.
13. Kunkov S, Pinedo V, Silver EJ, Crain EF. Predicting the need for hospitalization in acute childhood asthma using end-tidal capnography. *Pediatr Emerg Care*. 2005 Sep;21(9):574-7.
14. Lightdale JR, Goldmann DA, Feldman HA, Newburg AR, DiNardo JA, Fox VL. Microstream capnography improves patient monitoring during moderate sedation: a randomized, controlled trial. *Pediatrics* 2006 Jun;117(6):e1170-8.
15. Liu Z, Vargas F, Stansbury D, Sasse SA, Light RW. Comparison of the end-tidal arterial PCO₂ gradient during exercise in normal subjects and in patients with severe COPD. *Chest*. 1995 May;107(5):1218-24.
16. Looney JM, Jellinek EM. The oxygen and carbon dioxide content of the arterial and venous blood of normal volunteers. *Am J Physiol*. 1937; 118: 225-231
17. Miner JR, Heegaard W, Plummer D. End-tidal carbon dioxide monitoring during procedural sedation. *Acad Emerg Med*. 2002 Apr;9(4):275-80.
18. Repetto JE, Donohue PA-C PK, Baker SF, Kelly L, Noguee LM. Use of capnography in the delivery room for assessment of endotracheal tube placement. *J Perinatol*. 2001 Jul-Aug;21(5):284-7.
19. Sanchez O, Wermert D, Faisy C, Revel MP, Diehl JL, Sors H, Meyer G. Clinical probability and alveolar dead space measurement for suspected pulmonary embolism in patients with an abnormal D-dimer test result. *J Thromb Haemost*. 2006 Jul;4(7):1517-22.

20. Silvestri S, Ralls GA, Krauss B, Thundiyil J, Rothrock SG, Senn A, Carter E, Falk J. The effectiveness of out-of-hospital use of continuous end-tidal carbon dioxide monitoring on the rate of unrecognized misplaced intubation within a regional emergency medical services system. *Ann Emerg Med.* 2005 May;45(5):497-503.
21. Singh S, Allen WD Jr, Venkataraman ST, Bhende MS. Utility of a novel quantitative handheld microstream capnometer during transport of critically ill children. *Am J Emerg Med.* 2006 May;24(3):302-7.
22. Soto RG, Fu ES, Vila H Jr, Miguel RV. Capnography accurately detects apnea during monitored anesthesia care. *Anesth Analg.* 2004 Aug;99(2):379-82.
23. Sehra R, Underwood K, Checchia P. End tidal CO₂ is a quantitative measure of cardiac arrest. *Pacing Clin Electrophysiol* 2003 Jan;26(1 Pt 2):515-7
24. Vivien B, Amour J, Nicolas-Robin A, Vesque M, Langeron O, Coriat P, Riou B. An evaluation of capnography monitoring during the apnoea test in brain-dead patients. *Eur J Anaesthesiol.* 2007 Jun 20;:1-8
25. Yaron M, Padyk P, Hutsinpiller M, Cairns CB. Utility of the expiratory capnogram in the assessment of bronchospasm. *Ann Emerg Med.* 1996 Oct;28(4):403-7.
26. Yazigi A, Zeeni C, Richa F, Chalhoub V, Sleiaty G, Noun R. The accuracy of non-invasive nasal capnography in morbidly obese patients after bariatric surgery. *Middle East J Anesthesiol.* 2007 Oct;19(3):483-94.
27. Yosefy C, Hay E, Nasri Y, Magen E, Reisin L. End tidal carbon dioxide as a predictor of the arterial PCO₂ in the emergency department setting. *Emerg Med J.* 2004 Sep;21(5):557-9.
28. Zwerneman K. End-tidal carbon dioxide monitoring: a VITAL sign worth watching. *Crit Care Nurs Clin North Am.* 2006 Jun;18(2):217-25, xi

Pulse Oximetry

1. Ahrens TS, Tucker K. Pulse oximetry. *Critical Care Nursing Clinics of North America* 1999;11:87-98.
2. Giuliano KK, Liu LM. Knowledge of pulse oximetry among critical care nurses. *Dimens Crit Care Nurs* 2006 Jan-Feb;25(1):44-9.
3. Kuroda M, Kawamoto M, Yuge O. Undisrupted pulse wave on pulse oximeter display monitor at cardiac arrest in a surgical patient. *Journal of Anesthesia* 2005; 19:164 - 166
4. Ochroch EA, Russell MW, Hanson WC 3rd, et al. The impact of continuous pulse oximetry monitoring on intensive care unit admissions from a postsurgical care floor. *Anesth Analg* 2006 Mar;102(3):868-75.
5. Petterson MT, Begnoche VL, Graybeal JM. The effect of motion on pulse oximetry and its clinical significance. *Anesth Analg* 2007 Dec;105(6 Suppl):S78-84
6. Richards NM, Giuliano KK, Jones PG. A Prospective Comparison of 3 New-Generation Pulse Oximetry Devices During Ambulation After Open Heart Surgery. *Respir Care* 2006 Jan;51(1):29-35.
7. Severinghaus JW. Takuo Aoyagi: discovery of pulse oximetry. *Anesth Analg* 2007; Dec;105(6 Suppl):S1-4
8. Welch J. Pulse oximeters. *Biomed Instrum Technol.* 2005 Mar-Apr;39(2):125-30.
9. Workie FA, Rais-Bahrami K, Short BL. Clinical use of new-generation pulse oximeters in the neonatal intensive care unit. *Am J Perinatol* 2005 Oct;22(7):357-60.

Blood Gas Interpretation

1. Ahrens T, Prentice D, Kleinpell R. *Critical Care: Certification. Preparation and Review.* Norwalk: Appleton & Lange. 4th Ed. 2006.
2. Allen K. Four-step method of interpreting arterial blood gas analysis. *Nurs Times* 2005; Jan 4-10;101(1):42-5.
3. Shoulders-Odom B. Using an algorithm to interpret arterial blood gases. *Dimens Crit Care Nurs* 2000 Jan-Feb;19(1):36-41.
4. Simpson H. Interpretation of arterial blood gases: a clinical guide for nurses. *Br J Nurs* 2004 May 13-26;13(9):522-8.

